



A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie

GDCh

International Edition

www.angewandte.org

Accepted Article

Title: Optogenetic Control of Voltage Gated Calcium Channels

Authors: Yubin Zhou, Guolin Ma, Jindou Liu, Yuepeng Ke, Xin Liu, Minyong Li, Fen Wang, Gang Han, Yun Huang, and Youjun Wang

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Angew. Chem. Int. Ed.* 10.1002/anie.201713080
Angew. Chem. 10.1002/ange.201713080

Link to VoR: <http://dx.doi.org/10.1002/anie.201713080>
<http://dx.doi.org/10.1002/ange.201713080>

COMMUNICATION

Optogenetic Control of Voltage Gated Calcium Channels


Guolin Ma^{#[a]}, Jindou Liu^{#[b]}, Yuepeng Ke^{#[a]}, Xin Liu^[a], Minyong Li^[c], Fen Wang^[a], Gang Han^[d], Yun Huang^[a], Youjun Wang^{*[b]}, Yubin Zhou^{*[a,e]}

Abstract: Voltage-gated Ca²⁺ (Ca_v) channels mediate Ca²⁺ entry into excitable cells to regulate a myriad of cellular events following membrane depolarization. Here, we report the engineering of RGK GTPases, a class of genetically encoded Ca_v channel modulators, to enable photo-tunable modulation of Ca_v channel activity in excitable mammalian cells. This optogenetic toolkit (designated optoRGK) tailored for Ca_v channels will find broad applications in interrogating a wide range of Ca_v-mediated physiological processes in multiple systems.

Ca_v channels constitute the major route of Ca²⁺ entry in the nervous and cardiovascular systems, as well as other electrically excitable cells.^[1] Ca_v channels respond to membrane depolarization to permit Ca²⁺ influx, thereby playing instrumental roles in Ca²⁺-dependent physiological processes, including neurotransmitter/hormone release, gene expression and muscle contraction.^[1b, 1c] Deregulated Ca_v channels can give rise to pathophysiological conditions ranging from cardiovascular disorders to psychiatric conditions^[2]. Consequently, Ca_v channels are promulgated as important targets for therapeutic intervention and physiological regulation^[3]. Currently widely used Ca_v channel blockers (e.g., dihydropyridines) have prominent drawbacks, including off-target toxicity, lack of spatial control and non-reversibility.^[3b, 4] There remains, therefore, a salient need to exploit new interventional approaches to control Ca_v channels.

Optogenetics, which incorporates synthetic photosensitive modules into cells of living tissues to control cellular activities with high spatiotemporal precision, provides an ideal solution to

overcome the drawbacks associated with conventional Ca_v channel blockers.^[5] The Ras-like GTPases, Rad/Rem/Gem/Kir (RGK), which function as negative regulators of Ca_v channels, are considered as the prime candidates for generating optogenetic tools to modulate Ca_v channels.^[6] Given that membrane anchoring is necessary for RGKs to exert their suppressive effects on Ca_v channels,^[6b, 7] we reasoned that Ca_v channels could be remotely modulated by harnessing the power of light to control the translocation of RGK toward the plasma membrane (PM). We therefore engineered a set of optogenetic constructs by using a light-sensitive heterodimerization system to control the subcellular localization of engineered Rem (Scheme 1 and Figure S1). We chose the optical dimerizer pair made of iLID (LOV2-ssrA) and sspB because of their small sizes, fast photo responsive kinetics, low background interaction and minimized perturbation to endogenous signalling pathways.^[8]



Scheme 1. Design of optoRGK to photo-tune Ca_v channel activities. Spatiotemporal control of the Rem1 core domain is achieved by utilizing the LOV2-ssrA/sspB optical dimerizer. The light-inducible cytosol-to-PM translocation of Rem enables inducible suppression of Ca_v channel activity.

To enable light-inducible cytosol-to-PM translocation of engineered Rem, we set out to install sspB into different positions of mCherry tagged Rem₁₋₂₆₆ via flexible linkers with varying lengths (Figures S1-3). In parallel, we tethered Venus-iLID to PM with a C-terminal PM-targeting sequence (CAAX) derived from KRas4B^[8] or the PM-tethering motif (Lyn11) from the tyrosine-protein kinase Lyn.^[9] We then co-transfected these two constructs into HeLa cells and examined the reversible recruitment of cytosolic Rem₁₋₂₆₆ toward PM (Figure 1a, Figure S1-4). In the dark, mCherry-Rem₁₋₂₆₆-sspB was evenly distributed in the cytosol (Figure 1a). Upon blue light illumination, photoexcitation produced a covalent adduct between LOV2 residue C450 and the cofactor FMN, thereby exposing the caged ssrA component to restore its interaction with sspB. Consequently, mCherry-Rem₁₋₂₆₆-sspB was recruited toward PM within seconds ($t_{1/2, on} = 3.2 \pm 1.0$ sec). Upon withdrawal of light, mCherry-Rem₁₋₂₆₆-sspB dissociated from PM-

[a] Dr. G. Ma, Y. Ke, X. Liu, Dr. F. Wang, Prof. Y. Huang, Prof. Y. Zhou. Institute of Biosciences and Technology College of Medicine, Texas A&M University 2121 W Holcombe Blvd, Houston, TX 77030, USA E-mail: yzhou@ibt.tamhsc.edu

[b] J. Liu, Prof. Y. Wang Beijing Key Laboratory of Gene Resource and Molecular Development, College of Life Sciences, Beijing Normal University Beijing 100875, China E-mail: wyoujun@bnu.edu.cn

[c] Prof. M. Li, Department of Medicinal Chemistry, Key Laboratory of Chemical Biology, School of Pharmacy, Shandong University Jinan, Shandong 250012, China

[d] Prof. G. Hang, Department of Biochemistry and Molecular Pharmacology, University of Massachusetts Medical School Worcester, MA 01605

[e] Prof. Y. Zhou, Department of Medical Physiology, College of Medicine, Texas A&M University Temple, TX 76504, USA

[#] These authors contributed equally. Supporting information for this article is given via a link at the end of the document

COMMUNICATION

resident *ssrA* and diffused back to the cytosol ($t_{1/2, \text{off}} = 23.0 \pm 2.4$ sec) (Figure 1, S3 and Movie S1). After screening over 20 constructs with different combinations of key elements (various Rem fragments, linkers, PM-targeting motifs, and insertion sites), we identified mCherry-Rem₁₋₂₆₆-sspB-CAAX as an ideal candidate because it exhibited an optimal performance with high sensitivity and relatively fast kinetics (Figures S1-S3). To enable a near 1:1 co-expression of the two components using a single construct within the same cells, we resorted to a multicistronic expression system by utilizing the self-cleaving 2A-peptide (P2A)^[10]. We named the system as optoRGK and used this construct for further characterization and applications in excitable cells.




Figure 1. Visualization of the reversible recruitment of the Rem core domain (Rem₁₋₂₆₆ without the C-terminus) to PM in response to blue light illumination. **a**) Representative confocal images showing light-inducible cytosol-to-PM translocation of mCh-optoRGK in HeLa cells co-expressing Venus-iLID-CAAX. The images represent the same cell in the dark (black bar) or exposed to blue light at 470 nm (40 $\mu\text{W}/\text{mm}^2$; blue bar). Scale bar, 5 μm . **b**) Quantification of cytosolic mCherry signals of optoRGK over five repeated light-dark cycles. $n = 34$ cells from three independent experiments. Error bars denote s.e.m..

To determine whether optoRGK could photo-modulate Ca_v channels, we expressed optoRGK in HEK293 cells stably expressing the human $\text{Ca}_v1.2$ channel components (HEK- $\text{Ca}_v1.2$)^[11] and evaluated membrane depolarization induced Ca^{2+} entry with a red genetically encoded Ca^{2+} sensor jRCaMP1b^[12] (Figure 2a-b) or the green Ca^{2+} dye Fluo-4 (Figure S3). 50 mM potassium chloride (KCl) was employed to induce membrane depolarization. In the dark, addition of KCl elicited a pulse of Ca_v -mediated Ca^{2+} influx in both control and optoRGK-expressing cells (Figure 2, 1st cycle). Upon blue light illumination, cells overexpressing optoRGK showed a significant reduction in KCl-induced Ca^{2+} entry compared to control cells (Figure 2a, 2nd cycle). Furthermore, Ca^{2+} influx could be restored in the absence of blue light (Figure 2a, 3rd cycle), thereby attesting to the reversibility of optoRGK in the regulation of Ca_v channels. In parallel, we performed electrophysiological studies to independently confirm optoRGK-mediated, light-switchable modulation of Ca_v channel in HEK293- $\text{Ca}_v1.2$ cells. In the dark, cells expressing optoRGK

showed robust whole cell currents with a typical Ca_v I-V relationship and amplitudes similar to those of control cells (maximal peak current density: ~ 21.2 pA/pF, $n = 12$) (Figure 2c-d). However, upon receiving blue light stimulation, the amplitudes of depolarization-induced whole cell currents were significantly diminished (maximal peak current density: ~ 6.3 pA/pF, $n = 15$) (Figure 2c-d). These results clearly established optoRGK as a genetically encoded light-switchable channel modulator that allows optical inhibition of Ca_v channels in excitable cells.




Figure 2. OptoRGK-mediated photoswitchable inhibition of Ca^{2+} entry through $\text{Ca}_v1.2$ channels. **a**) Ca^{2+} influx in HEK- $\text{Ca}_v1.2$ cells transiently expressing optoRGK and a red Ca^{2+} sensor jRCaMP1b with and without blue light stimulation. Cells transfected with the empty vector is used as control. Membrane depolarization induced Ca^{2+} entry was elicited by adding 50 mM KCl (black line below the curves; three repeated cycles) to transfected cells. Blue line represents light stimulation under 470 nm with a power density of 40 $\mu\text{W}/\text{mm}^2$. **b**) Bar graphs showing the statistical results of mean Ca^{2+} entry for each cycle. **c**) The current-voltage relationships of Ca_v channels in HEK- $\text{Ca}_v1.2$ cells transfected with optoRGK. Cells were either shielded from light or exposed to blue light prior to electrophysiological recording. **d**) Bar graphs showing the statistical results of peak whole cell currents induced by pulses of +10 mV depolarization (pA/pF) in HEK- $\text{Ca}_v1.2$ cells before and after photo-stimulation. All data were presented as mean \pm s.e.m. $**P < 0.01$ (paired Student's *t*-test).

We next moved on to test optoRGK in C2C12 cells, a mouse myoblast cell line^[13] with functional Ca_v channels^[6b]. Again, we observed light-dependent inhibition of Ca^{2+} influx in this excitable cell line (Figure S5). To further test optoRGK in more physiologically-relevant systems, we introduced it into HL-1 cardiomyocytes, a well-characterized atrial myocyte culture line derived from the adult mouse that retains many of the differentiated properties of cardiac cells^[14], including rhythmic oscillations of cytosolic Ca^{2+} (Movie S2). By using Fluo-4 as the Ca^{2+} indicator, we first evaluated rhythmic Ca^{2+} oscillations in HL-1 cells with and without overexpression of full length Rem or its truncated version, Rem₁₋₂₆₆ (Figure S6, Movies S3 and S4). Both control cells and HL-1 cells transfected with mCh-Rem₁₋₂₆₆ (Figure S6b) exhibited rhythmic Ca^{2+} oscillations, while cardiomyocytes expressing the full-length Rem failed to evoke Ca^{2+} transients

COMMUNICATION

(Figure S6a). These findings were concordant with the results obtained from C2C12 myoblast cells (Figure S5) and other types of excitable cells^[15].




Figure 3. OptoRGK-mediated light-inducible inhibition of the rhythmic oscillations of cytosolic Ca^{2+} in cardiac cells. **a)** Ca^{2+} oscillations reported by Fluo-4 in HL-1 cells with (bottom) and without (Top) expression of optoRGK. Kymographs of Fluo-4 signals in a representative HL-1 cell were shown above the traces. Excitation was set at 488 nm to record Fluo-4 signals while simultaneously photoactivating optoRGK. **b)** Ca^{2+} oscillations in HL-1 cells monitored by Cal-590. Blue bar, blue light illumination at 470 nm ($40 \mu\text{W}/\text{mm}^2$).

Having validated the use of HL-1 cardiomyocytes to test our tool, we next examined the rhythmic oscillations of cytosolic Ca^{2+} (Fluo-4 signals as readout) in HL-1 cells expressing optoRGK. Upon blue light illumination, mCh-Rem₁₋₂₆₆ translocated from the cytosol to close to PM within several seconds (Figure S7), accompanied by the attenuation of oscillatory Ca^{2+} signals (bottom, Figure 3a, Movie S5). By contrast, the control cells displayed regular Ca^{2+} oscillations under blue light. To further validate if such action was reversible in HL-1 cells, we used a red Ca^{2+} indicator, Cal-590 (excitation at 562 nm without pre-activating optoRGK), rather than Fluo-4 to monitor Ca^{2+} oscillations (Figure 3b). HL-1 cells expressing optoRGK showed regular Ca^{2+} spikes in the dark. However, upon blue light illumination, the rhythmic oscillations were substantially attenuated. Notably, regular Ca^{2+} oscillations were restored in the same HL-1 cardiomyocyte after removal of the light source (Figure 3b).

Taken together, compared with traditional small molecule Ca_v channel blockers that often lack reversibility, selectivity and tissue-specificity, engineered RGK proteins could serve as promising candidates to enable spatiotemporal control of Ca_v channels with a simple pulse of light. This study complements the recent development of engineered stromal interaction molecule 1 (STIM1) to photo-regulate endogenous Ca^{2+} channels in mammalian cells (e.g., optoSTIM^[16] and Opto-CRAC^[17]). We anticipate that the optoRGK toolkit developed in the current study will find broad applications in interrogating a wide range of Ca^{2+} -dependent physiological processes in mammals.

Proof-of-concept experiments have already demonstrated the potential of using RGK to treat heart disease^[18]. To test potential in vivo applications, it is our immediate future plan to express

optoRGK in the atrioventricular node of rodent models with atrial fibrillation disease^[19], and examine whether photostimulation could suppress aberrant atrioventricular nodal conduction to intervene atrial fibrillation.

Given that RGK proteins primarily exert suppressive effects on high voltage-activated Ca^{2+} ($\text{Ca}_v1/\text{Ca}_v2$) channels^[20] and that LOV2-based photoswitches have a relatively slow kinetics, the efficacy of optoRGK will likely depend on the distribution and endogenous levels of Ca_v channel subtypes in different cell types and tissues. At its current configuration, optoRGK is well suited to modulate cardiomyocytes because of the abundant expression of L-type $\text{Ca}_v1.1/\text{Ca}_v1.2$ channels and the relatively long duration of cardiac action potentials. Its compatibility with neurons and other types of electrically excitable cells (e.g., pancreatic beta cells), remain to be rigorously tested in future studies.

Acknowledgements

This work was supported by grants from NIH (R21GM126532 and R01GM112003), CPRIT (RP170660 and RR140053), the National Natural Science foundation of China (NSFC-31471279 and NSFC-81728019), the Recruitment Program for Young Professionals of China, the Fundamental Research Funds for the Central Universities (2017STUD20), the Welch Foundation (BE-1913) and the American Cancer Society (RSG-16-215-01-TBE).

Keywords: ion channels • optogenetics • calcium signalling • protein engineering • synthetic biology

- [1] aS. N. Yang, P. O. Berggren, *American journal of physiology. Endocrinology and metabolism* **2005**, 288, E16-28; bD. Atlas, *Annual review of biochemistry* **2013**, 82, 607-635; cW. A. Catterall, *Annual review of cell and developmental biology* **2000**, 16, 521-555.
- [2] aI. Splawski, K. W. Timothy, L. M. Sharpe, N. Decher, P. Kumar, R. Bloise, C. Napolitano, P. J. Schwartz, R. M. Joseph, K. Condouris, H. Tager-Flusberg, S. G. Priori, M. C. Sanguinetti, M. T. Keating, *Cell* **2004**, 119, 19-31; bB. A. Simms, G. W. Zamponi, *Neuron* **2014**, 82, 24-45.
- [3] aG. W. Zamponi, J. Striessnig, A. Koschak, A. C. Dolphin, *Pharmacological reviews* **2015**, 67, 821-870; bG. W. Zamponi, *Nature reviews. Drug discovery* **2016**, 15, 19-34.
- [4] C. E. Sunkel, M. Fau de Casa-Juana, F. J. Cillero, J. G. Priego, M. P. Ortega, *Journal of medicinal chemistry* **1988**, 31, 1886-1890.
- [5] M. Wang, Y. Yu, J. Shao, B. C. Heng, H. Ye, *Quantitative Biology* **2017**, 1-13.
- [6] aP. Beguin, K. Nagashima, T. Gono, T. Shibusaki, K. Takahashi, Y. Kashima, N. Ozaki, K. Geering, T. Iwanaga, S. Seino, *Nature* **2001**, 411, 701-706; bB. S. Finlin, S. M. Crump, J. Satin, D. A. Andres, *Proceedings of the National Academy of Sciences of the United States of America* **2003**, 100, 14469-14474.
- [7] aR. N. Correll, C. Pang, B. S. Finlin, A. M. Dailey, J. Satin, D. A. Andres, *The Journal of biological chemistry* **2007**, 282, 28431-28440; bT. Yang, Y. Suhail, S. Dalton, T. Kernan, H. M. Colecraft, *Nature chemical biology* **2007**, 3, 795-804.
- [8] G. Guntas, R. A. Hallett, S. P. Zimmerman, T. Williams, H. Yumerefendi, J. E. Bear, B. Kuhlman, *Proceedings of the National Academy of Sciences of the United States of America* **2015**, 112, 112-117.
- [9] G. R. Hammond, M. J. Fischer, K. E. Anderson, J. Holdich, A. Koteci, T. Balla, R. F. Irvine, *Science* **2012**, 337, 727-730.

COMMUNICATION


- [10] J. H. Kim, S. R. Lee, L. H. Li, H. J. Park, J. H. Park, K. Y. Lee, M. K. Kim, B. A. Shin, S. Y. Choi, *PLoS one* **2011**, *6*, e18556.
- [11] B. Balasubramanian, J. P. Imredy, D. Kim, J. Penniman, A. Lagrutta, J. J. Salata, *Journal of pharmacological and toxicological methods* **2009**, *59*, 62-72.
- [12] H. Dana, B. Mohar, Y. Sun, S. Narayan, A. Gordus, J. P. Hasseman, G. Tsegaye, G. T. Holt, A. Hu, D. Walpita, R. Patel, J. J. Macklin, C. I. Bargmann, M. B. Ahrens, E. R. Schreiter, V. Jayaraman, L. L. Looger, K. Svoboda, D. S. Kim, *eLife* **2016**, *5*.
- [13] S. Burattini, P. Ferri, M. Battistelli, R. Curci, F. Luchetti, E. Falcieri, *European journal of histochemistry : EJH* **2004**, *48*, 223-233.
- [14] aW. C. Claycomb, N. A. Lanson, Jr., B. S. Stallworth, D. B. Egeland, J. B. Delcarpio, A. Bahinski, N. J. Izzo, Jr., *Proceedings of the National Academy of Sciences of the United States of America* **1998**, *95*, 2979-2984; bL. Sartiani, P. Bochet, E. Cerbai, A. Mugelli, R. Fischmeister, *The Journal of physiology* **2002**, *545*, 81-92.
- [15] aD. D. Chang, H. M. Colecraft, *The Journal of physiology* **2015**, *593*, 5075-5090; bJ. Magyar, C. E. Kiper, G. Sievert, W. Cai, G. X. Shi, S. M. Crump, L. Li, S. Niederer, N. Smith, D. A. Andres, J. Satin, *Channels* **2012**, *6*, 166-173.
- [16] aG. Ma, S. Wen, L. He, Y. Huang, Y. Wang, Y. Zhou, *Cell calcium* **2017**, *64*, 36-46; bT. Kyung, S. Lee, J. E. Kim, T. Cho, H. Park, Y. M. Jeong, D. Kim, A. Shin, S. Kim, J. Baek, J. Kim, N. Y. Kim, D. Woo, S. Chae, C. H. Kim, H. S. Shin, Y. M. Han, D. Kim, W. D. Heo, *Nature biotechnology* **2015**, *33*, 1092-1096.
- [17] L. He, Y. Zhang, G. Ma, P. Tan, Z. Li, S. Zang, X. Wu, J. Jing, S. Fang, L. Zhou, Y. Wang, Y. Huang, P. G. Hogan, G. Han, Y. Zhou, *eLife* **2015**, *4*.
- [18] aC. A. Makarewich, R. N. Correll, H. Gao, H. Zhang, B. Yang, R. M. Berretta, V. Rizzo, J. D. Molkenin, S. R. Houser, *Circulation research* **2012**, *110*, 669-674; bM. Murata, E. Cingolani, A. D. McDonald, J. K. Donahue, E. Marban, *Circulation research* **2004**, *95*, 398-405.
- [19] K. Nishida, G. Michael, D. Dobrev, S. Nattel, *Europace : European pacing, arrhythmias, and cardiac electrophysiology : journal of the working groups on cardiac pacing, arrhythmias, and cardiac cellular electrophysiology of the European Society of Cardiology* **2010**, *12*, 160-172.
- [20] T. Yang, H. M. Colecraft, *Biochimica et biophysica acta* **2013**, *1828*, 1644-1654.

COMMUNICATION

Entry for the Table of Contents (Please choose one layout)

COMMUNICATION

We report herein the invention of optoRGK, a new class of engineered modulators of voltage-gated calcium (Ca_v) channels, to enable photo-tunable regulation of Ca_v channel activity and Ca^{2+} entry in excitable mammalian cells, such as cardiomyocytes.



Guolin Ma, Jindou Liu, Yuepeng Ke, Xin Li, Minyong Li, Fen Wang, Gang Han, Yun Huang, Youjun Wang, Yubin Zhou

Page No. – Page No.

Optogenetic Control of Voltage Gated Calcium Channels